

UCSF

UC San Francisco Previously Published Works

Title

Draft Genome Sequence of Mycobacterium elephantis Strain Lipa.

Permalink

<https://escholarship.org/uc/item/0d11k579>

Journal

Genome announcements, 3(3)

ISSN

2169-8287

Authors

Greninger, Alexander L
Cunningham, Gail
Yu, Joanna M
et al.

Publication Date

2015-06-01

DOI

10.1128/genomea.00691-15

Peer reviewed

Draft Genome Sequence of *Mycobacterium elephantis* Strain Lipa

Alexander L. Greninger, Gail Cunningham, Joanna M. Yu, Elaine D. Hsu, Charles Y. Chiu, Steve Miller

Department of Laboratory Medicine, UCSF, San Francisco, California, USA

We report the draft genome sequence of *Mycobacterium elephantis* strain Lipa from a sputum sample of a patient with pulmonary disease. This is the first draft genome sequence of *M. elephantis*, a rapidly growing mycobacterium.

Received 20 May 2015 Accepted 22 May 2015 Published 25 June 2015

Citation Greninger AL, Cunningham G, Yu JM, Hsu ED, Chiu CY, Miller S. 2015. Draft genome sequence of *Mycobacterium elephantis* strain Lipa. *Genome Announc* 3(3):e00691-15. doi:10.1128/genomeA.00691-15.

Copyright © 2015 Greninger et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Steve Miller, steve.miller@ucsf.edu.

Mycobacterium elephantis is a rapidly growing mycobacterium, originally isolated from a lung abscess in an elephant in 2000, that has been occasionally isolated from human clinical samples, most commonly sputum specimens (1–3). *M. elephantis* has also been reported to have been isolated from a cervical lymph node, although its relation to clinical disease is unclear (2). The notable phenotypic characteristics of *M. elephantis* include a relatively slow growth rate for a rapid growing mycobacterium, smooth colonies producing a pale yellow pigment with age, and ability to grow on 5% NaCl in Lowenstein-Jensen medium, along with a unique high-performance liquid chromatography (HPLC) mycolic acid profile (3).

Rapidly growing mycobacteria constitute a commonly isolated population of acid-fast bacillus in the clinical microbiology lab of varying clinical importance (4, 5). We sequenced the first draft genome of *M. elephantis* from a sputum sample of a patient in 2003 with pulmonary disease. The isolate was originally typed as *M. elephantis* based on partial 16S sequencing.

DNA from *M. elephantis* strain Lipa was extracted using the Qiagen EZ1 kit, and paired-end libraries were prepared using the Nextera XT DNA library kit followed by sequencing on an Illumina MiSeq instrument. Sequences were adapter and quality (Q20) trimmed using Cutadapt, *de novo* assembled using SPAdes v3.5, metagenomically screened for contaminating sequence with SURPI, and annotated via Prokka v1.1 (6–9). A total of 9,000,614 paired-end reads of average length 115 nucleotides were recovered after trimming. *De novo* assembly yielded 234 contigs for a total assembly size of 5,187,616 bp with an N_{50} of 42,430 bp, an average coverage of 196×, and a total of 5,022 coding sequences. Contiguity was most likely disrupted by the high G+C content (68%) along with several high-copy-number integrases, transposases, and recombinases that were longer than sequence read length. Other high-copy-number contigs included those containing genes to *mleA* phospholipid ABC transporter permease, for which *M. elephantis* strain Lipa had 19 different homologs in the genome.

The five closest BLASTN hits to the complete 16S sequence from the isolate were other *M. elephantis* 16S sequences with 99.4 to 100% identity. The Lipa strain contained three putative beta-glucosidase genes. By Comprehensive Antibiotic Resistance Database analysis, the Lipa strain includes an *arr1* rifampin ADP-

ribosyl transferase (85% by amino acid to *M. vanbaalenii* PYR-1) and a *blaF* beta-lactamase (10).

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession no. [LBNO000000000](https://www.ncbi.nlm.nih.gov/nuclseq/LBNO000000000). The assembly described in this paper is the first version, LBNO01000000.

ACKNOWLEDGMENT

This work was supported by the UCSF Department of Laboratory Medicine.

REFERENCES

- Shojaei H, Magee JG, Freeman R, Yates M, Horadagoda NU, Goodfellow M. 2000. *Mycobacterium elephantis* sp. nov., a rapidly growing non-chromogenic mycobacterium isolated from an elephant. *Int J Syst Evol Microbiol* 50:1817–1820.
- Turenne C, Chedore P, Wolfe J, Jamieson F, May K, Kabani A. 2002. Phenotypic and molecular characterization of clinical isolates of *Mycobacterium elephantis* from human specimens. *J Clin Microbiol* 40:1230–1236. <http://dx.doi.org/10.1128/JCM.40.4.1230-1236.2002>.
- Tortoli E, Rindi L, Bartoloni A, Garzelli C, Mantella A, Mazzarelli G, Piccoli P, Scarparo C. 2003. *Mycobacterium elephantis*: not an exceptional finding in clinical specimens. *Eur J Clin Microbiol Infect Dis* 22:427–430. <http://dx.doi.org/10.1007/s10096-003-0950-2>.
- Greninger AL, Langelier C, Cunningham G, Keh C, Melgar M, Chiu CY, Miller S. 29 April 2015. Two rapidly growing mycobacterial species isolated from a brain abscess: first whole genome sequence of *Mycobacterium immunogenium* and *Mycobacterium llutzerense*. *J Clin Microbiol*. <http://dx.doi.org/10.1128/JCM.00402-15>.
- De Groote MA, Huitt G. 2006. Infections due to rapidly growing mycobacteria. *Clin Infect Dis* 42:1756–1763. <http://dx.doi.org/10.1086/504381>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
- Naccache SN, Federman S, Veeraraghavan N, Zaharia M, Lee D, Samayoa E, Bouquet J, Greninger AL, Luk K-C, Enge B, Wadford DA, Messenger SL, Genrich GL, Pellegrino K, Grard G, Leroy E, Schneider BS, Fair JN, Martinez MA, Isa P, Crump JA, DeRisi JL, Sittler T, Hackett J, Miller S, Chiu CY. 2014. A cloud-compatible bioinformatics pipeline for ultrarapid pathogen identification from next-generation sequencing of clinical samples. *Genome Res* 24:1180–1192. <http://dx.doi.org/10.1101/gr.171934.113>.
- Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17:10–12. <http://dx.doi.org/10.14806/ej.17.1.200>.
- Bankovich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov

- AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
10. McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, De Pascale G, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJ, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang W, Yan M, Yu T, Wright GD. 2013. The comprehensive antibiotic resistance database. *Antimicrob Agents Chemother* 57:3348–3357. <http://dx.doi.org/10.1128/AAC.00419-13>.